

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Role Of CT Urography In Investigating Hematuria.

Mohamed Riaz*.

Assistant Professor, Department Of Radio Diagnosis, Government Medical College And Hospital, Ariyalur Tamil Nadu, India.

ABSTRACT

Hematuria can signify serious diseases such as bladder cancer, upper urinary tract urothelial cell cancer (UUT-UCC), renal cell cancer, or urinary tract stones. To analyze the main role of CT urography in case of hematuria in terms of diagnosis and treatment. This cross-sectional study was conducted in the Department of Radiology Government Medical College And Hospital, Ariyalur Tamil Nadu, India from January 2024 to June 2024. The data was collected from 50 patients of haematuria from the OPD of the hospital. The patients underwent a CECT examination after obtaining a detailed clinical history. There were 38 females and 12 males. The most common cause of obstructive uropathy was stone disease i.e. renal, ureteric or both and 75.0% patients in group A and 65.0% in group B, presented with it followed by other causes i.e. carcinomas, pyonephrosis, and posterior urethral jugular vein (PUJ) obstruction. It is concluded that multidetector CT urography detects the entire spectrum of urinary tract pathologies causing haematuria with high accuracy.

Keywords: Clinical Study, CT, Urography, Hematuria

<https://doi.org/10.33887/rjpbcs/2025.16.2.17>

**Corresponding author*

INTRODUCTION

In some cases, hematuria can indicate the presence of bladder cancer, UUT-UCC, renal cell carcinoma, or urinary tract stones. The presence of blood in the urine, whether microscopic or macroscopic, is referred to as haematuria. More precisely, it is defined as the presence of five or more red blood cells that were collected at least one week apart from one another. The presence of blood in the urine is usually the first sign of hematuria in the clinical setting. Symptomatic or asymptomatic, short-term or long-term, isolated or in conjunction with other urinary abnormalities, it can occur as a single finding or as part of a larger pattern [1]. As one of the most common urinary tract pathologies, haematuria warrants both the patient's and treating physician's attention. One of the most common symptoms of a variety of urinary tract disorders, including calculi, neoplasms, infections, trauma, medication-induced thrombocytopenia, developmental anomalies, and diseases of the renal parenchyma, is haematuria [2]. Intravenous urography (IVU) was the first method of genitourinary imaging until the beginning of the 21st century. However, MDCT urography is now the preferred imaging technique [3]. Haematuria investigation depends on practice however, this has changed recently. Plain film imaging may be useful in certain situations, such as in the case of acute renal colic that results in haematuria. To rule out malignancy, the most sensitive imaging modalities are ultrasound and CT [4]. Detection of haematuria relies heavily on the use of ultrasound. Its sensitivity varies based on the operator's experience and the patient's body habitus. Urinary tract calculi can be detected using this method. Further research shows that ultrasound is more sensitive than IVU for the detection of urological malignancy [5]. When it comes to diagnosing haematuria, contrast-enhanced CT is most effective. Compared to ultrasound's sensitivity of about 85% in diagnosing renal masses, its sensitivity ranges from >98% to >85% for this test. By contrast, 15 CT-U has an 85 percent higher sensitivity in diagnosing upper tract urothelial malignancy than IVU [6].

METHODS

This cross-sectional study was conducted in the Department of Radiology Government Medical College And Hospital, Ariyalur Tamil Nadu, India from January 2024 to June 2024. The data was collected from 50 patients of haematuria from the OPD of the hospital. The patients underwent a CECT examination after obtaining a detailed clinical history. Fifty patients with haematuria from the hospital's OPD were included. This study included all patients with haematuria between the ages of 30 and 60 years. Excluded from the study were patients with cardiovascular and renal disease, as well as those who were unwilling to participate. After obtaining a thorough clinical history, the patients were subjected to a CECT examination. Six hours before the study, the patient is instructed to fast. Non-contrast phase: First, the non-contrast stage, and then the corticomedullary phase, which was acquired following a delay of 25-80 seconds after administration of non-ionic low osmolar contrast medium to distinguish normal variants of renal parenchymal from renal masses and better depiction tumor hypervascularity, respectively. CT scans were taken from the diaphragm to the bladder using a Multi-detector row CT scanner. Histopathological findings were used to determine the next step in the diagnosis process. SPSS version 20 was used to collect and analyze the data. The mean and standard deviation were used for all of the data.

RESULTS

A total of 50 patients participated in the study. Thirty-eight women and twelve men took part. Urinary obstruction was caused most frequently by renal, ureteric, or both stone disease, which was found in 75.0% of group A patients and 65.0% of those in group B, as shown in Table 1. Other common causes included carcinomas, pyonephrosis, and obstruction of the posterior urethral jugular vein (PUJ) (Table 2).

Table 1: Causes of Obstructive Uropathy in selected patients

Causes	% Age
Stone disease	75.0
Renal	40.0
Ureteric	25.0
Renal + Ureteric	10.0
Carcinomas	20.0
Urinary Bladder	3.0

Prostate	2.0
Cervix	5.010.0
Others	
Pyonephrosis	3.0
PUJ Obstruction	2.0

Cervical cancer was the most common malignancy associated with ureteral obstructions requiring urinary diversion among the benign etiologies.

Table 2: Primary cause of ureteral obstruction

Causes	Ureteral stent	Percutaneous nephrostomy
Benign cause	40	16
Malignancy	26	28
Cervical cancer	19	9
Prostate cancer	4	5
colon Cancer	1	7
Bladder Cancer	2	1
Stomach cancer	0	1
Ovarian cancerate	0	1
lung cancer	0	1
Endometrial cancer	0	1
Lymphoma	0	1
Breast Cancer	0	1

A total of 24 of the 27 neoplastic foci on MDCT urography were found by the two retrospective reviewers, including all 18 prospectively discovered urothelial tumors as well as six additional lesions. A total of six patients were diagnosed with these abnormalities. Two malignancies that were missed in the prospective review were found retroactively in four patients. Three small masses in the intrarenal collecting system constituted the six foci that were not seen in advance but were later discovered (Figure 1).

Figure 1: CT of a patients shows a soft tissue defect



DISCUSSION

Contrast-opacified thin section images of the collecting systems, ureters, and bladder can be added to conventional abdominal CT scans to enhance the examination of the entire urinary tract, which is known as CT urography [8]. When intravenous contrast, previously administered for the CT, has rendered the urinary tract opaque, film (or direct digital) images can be exposed to obtain the intravenous- pyelogram-like portions of the examination. These images can also be created by reformatting delayed CT scans in the axial or transverse planes [9]. A pyelogram portion of this exam may be comparable to a standard intravenous pyelogram, making CT more sensitive and specific (statistically

and pathologically) than ultrasound or nephrotomography about focal renal parenchymal abnormalities [10-12]. The urinary system has recently been evaluated with MRI. An inherent drawback to this modality's utility in diagnosing urinary pathologies, however, is that MRI cannot detect calcination. Its use is also constrained by the high price and difficulty in obtaining it [13]. For the time being, the use of MR urography is restricted to pregnant women, children, those with renal insufficiency, and eight patients who are allergic to contrast agents. To better understand urinary tract disorders associated with hemorrhagic cystitis, many authors have proposed using CT urography to assess both the renal parenchyma and urothelium [14]. Every patient who undergoes a CT urogram receives water, which is primarily used to hydrate and expand the kidneys and urinary tract. It is next necessary to perform non-contrast helical CT scan of the kidney to detect renal calculi [15]. To check for tumors and filling defects, a high-resolution nephrographic phase (slices of 1 to 2 mm thick) is performed, followed by an injection of iodinated contrast media [16,17]. The latter can be used to reconstruct the urinary system and the bladder for diagnostic purposes. When combined with delayed images similar to those from an intravenous- pyelogram, this type of CT urography is just as sensitive as the more traditional intravenous pyelogram in diagnosing the underlying cause of hematuria [18-20]. This study didn't take into account other risk factors for cancer, such as age. A patient's history of bladder cancer or gross rather than microscopic hemorrhage was cited in some examination reports, but risk factors were not specified in the vast majority of patient histories or medical charts. Patients with gross hematuria were more likely than those with microscopic hematuria to be diagnosed with malignancy.

CONCLUSION

It is concluded that Multiple-detector CT urography can accurately detect the full range of urinary tract pathologies that cause haematuria. Urinary tract pathologies can be accurately diagnosed using CT Urography.

REFERENCES

- [1] Alderson SM, Hilton S, Papanicolaou N. CT urography: Review of technique and spectrum of diseases. *Applied Radiology* 2011; 40(7):6.
- [2] Bhatt S, Dogra VS. *Kidney In: CT and MRI of the Whole Body*. 5th ed. Philadelphia, Pa: Mosby. 2008; 2:1863 - 1952
- [3] Cancers in Patients Evaluated for Microhematuria: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2021; 4(5): e218409.
- [4] Feroze S, Singh B, Gojwari T, Manjeet S, Athar B, Hamid H. Role of non-contrast spiral computerized tomography in acute ureteric colic. *Indian Journal of Urology: U: Journal of the Urological Society of India* 2007; 23(2):119.
- [5] Jinzaki M, Tanimoto A, Shinmoto H, Horiguchi Y, Sato K, Kuribayashi S, Silverman SG. Detection of bladder tumors with dynamic contrast-enhanced MDCT. *American Journal of Roentgenology* 2007;188(4):9138.
- [6] Lin MY, Li WM, Huang CN, Lee HL, Niu SW, Chen LT, et al. Dialysis increases the risk of bladder recurrence in patients with upper tract urothelial cancer: A population-based study. *Ann Surg Oncol* 2018; 25:1- 8.
- [7] Maher MM et al, Urinary Tract Imaging with Multidetector CT Urography in Patients with Hematuria. *Korean Journal of Radiology*, 2004; 5(1). Fielding JR, Fox LA, Heller H, Seltzer SE, Tempany CM, Silverman SG et al . In the evaluation of flank pain: overall accuracy and feature analysis. *Journal Of Computer-Assisted Tomography* 1997; 21(4):635 -8.
- [8] Maher MM, Kalra MK, Rizzo S, Mueller PR, Saini S. Multidetector CT urography in imaging of the urinary tract in patients with hematuria. *Korean Journal Of Radiology* 2004;5(1):1-0.
- [9] Owan NC, Turney BW, Taylor NJ, McCarthy CL, Crew JP. Multidetector computed tomography urography for diagnosing upper urinary tract urothelial tumor. *BJU international* 2007;99(6):1363-70.
- [10] Rud, E., Galtung, K.F., Lauritzen, P.M. et al. Examining upper urinary tract In patients with hematuria—time to revise the CT urography protocol? *Eur Radiol* 2020; 30, 1664–1670.
- [11] Silverman SG, Leyendecker JR, Amis Jr ES. What is the current role of CT urography and MR urography in the evaluation of the urinary tract? *Radiology*. 2009; 250(2):309 -23.
- [12] Song JH ,Beland MD, Mayo-Smith WW.Hematuria evaluation with MDCT urography.*AJR* 2011 ;197:W84- W89
- [13] Van Der Molen A, et al. CT urography: definition, indications and techniques. A guideline for



- clinical practice. *European Radiology* 2018;18, 4-17.
- [14] Yap LC, et al. Upper tract imaging modality to investigate haematuria: cancer detection rates and changing guidelines. *Afr J Urol* 2021; 27, 112
- [15] Yoo S, You D, Song C, Hong B, Hong JH, Kim CS, et al. Risk of intravesical recurrence after ureteroscopic biopsy for upper tract urothelial carcinoma: Does the location matter? *J Endourol* 2017; 31:259-65
- Ketata S, Ketata H, Sahnoun A, Slimen MH, Fakhfakh H, Bahloul A et al. Laterality of symptomatic recurrent calcium nephrolithiasis. *African Journal of Urology*. 2008; 14(3):143 -6